

VOLUME 2, ISSUE 1

Journal of ongoing Chemical Research

BOERHAVIA AND ITS PHYTOCHEMISTRY

Said Nadeem

APRIL 04, 2015

Nyctagenaceae is a family of 30 genera and c. 290 species in tropical and subtropical parts of both the hemisphere especially in America. In Pakistan it is represented by 5 genera and 10 or 11 species. Plants are trees, herbs. Some plants are ornamental e.g. *Mirabilis jalapa*. *Boerhavia* (*Boerhaavia*) is genus of this family with c. 40 species. Plants are mostly herbs with opposite leaves, found in tropical and subtropical regions, S. Europe, and S. United States. In Pakistan it is represented by 4 or 5 species. Anthocarp of *Boerhavia* is 5-ribbed, which is the main difference in the *Boerhavia* and *Commicarpus*, having 10-ribbed anthocarp. A mini review is presented which cover data upto 2008.

BOERHAVIA AND ITS PHYTOCHEMISTRY

Said Nadeem,^{a,b} Mehmet A. Özler,^{a,b} Mehmet Öztürk,^a Mehmet E. Duru,^a

^a Faculty of Sciences, Department of Chemistry, Muğla Sıtkı Koçman University, Kötekli-48121, Muğla Turkey

^b Department of Medicinal and Aromatic Plants, Koyceyiz Vocational School, Muğla Sıtkı Koçman University, Koyceyiz-48000 Muğla, Turkey

*Corresponding Author Email: saidnadeem@mu.edu.tr

Phone: +905335499880


ABSTRACT


Nyctagenaceae is a family of 30 genera and c. 290 species in tropical and subtropical parts of both the hemisphere especially in America. In Pakistan it is represented by 5 genera and 10 or 11 species. Plants are trees, herbs. Some plants are ornamental e.g. *Mirabilis jalapa*. *Boerhavia* (*Boerhaavia*) is genus of this family with c. 40 species. Plants are mostly herbs with opposite leaves, found in tropical and subtropical regions, S. Europe, and S. United States. In Pakistan it is represented by 4 or 5 species. Anthocarp of *Boerhavia* is 5-ribbed, which is the main difference in the *Boerhavia* and *Commicarpus*, having 10-ribbed anthocarp. A mini review is presented which cover data upto 2008.


INTRODUCTION


Nyctagenaceae is a family of 30 genera and c. 290 species in tropical and subtropical parts of both the hemisphere especially in America. In Pakistan it is represented by 5 genera and 10 or 11 species. Plants are trees, herbs. Some plants are ornamental e.g. *Mirabilis jalapa*. *Boerhavia* (*Boerhaavia*) is genus of this family with c. 40 species. Plants are mostly herbs with opposite leaves, found in tropical and subtropical regions, S. Europe, and S. United States. In Pakistan it is represented by 4 or 5 species. Anthocarp of *Boerhavia* is 5-ribbed, which is the main difference in the *Boerhavia* and *Commicarpus*, having 10-ribbed anthocarp [3]. Recently nutritive and antinutritive properties of *B. difussa* has been published [35]. Hypoglycaemic effect of the aqueous extract of *B. diffusa* has been reported by Chude et al [36]. *Diffusa* leavesboerhavia *diffusa* is of great medicinal and ayurvedic importance. Red variety of *B. diffusa* is used as astringent to the bowel, used in biliousness, blood impurities, leucorrhea, anemia, and inflammation. Leaves of dark variety are used in dyspepsia, tumors, and enlargement of the spleen, abdominal pains, ophthalma, eyes wounds, and pain of the joints. Seeds are tonic, expectorant, carminative, muscular pain, lumbago, scabies, scorpion-sting, purify blood, hasten and delivery. Root has diuretic properties and also good expectorant, jaundice, ascites, anasarca, scanty urine internal inflammations, laxative and stomachic. When mixed with dried ginger, used in urticaria. The extract is used in dropsical swellings, gonorrhea, snake-bite, strangury, emetic, cardiac stimulant, myalgia, constipation, bronchitis, and cough [2]. Plants are also used as pot-herb with succulent leaves, fodder for cattle. The leaves are given to hogs in Jamaica, in Bengal, given to cattle to increase the milk quantity [1]. A large number of compounds belonging to flavonoids, alkaloids, carbohydrates, lipids, sterols, cyanine etc have been reported from the plants of this genus which are given in the table 1. The phytochemical studies on genus *Boerhavia* trace back to 1947, when Basu [34] et al first isolated punarnavine, an alkaloid. In 1954, Lightelm reported the presence of triacontenoic acid (**63**) in *B. repens* [29]. In 1968, T.N. Bhalla and M.L. Gupta reported the presence of punarnavine, an alkaloid of great medicinal importance, from *B. diffusa* [28]. In the same year Ahmad et al [27] isolated Hypoxanthine-9-L-arabinofuranoside (**28**) from the same plant. Hentriacontane (**65**) was

reported by A.N. Misra and H.P. Tiwari in 1971. After three year later, Khalique et al reported the presence of myristic acid and myricyl alcohol from *B.repens* [26]. Myristic alcohol was also reported by Suri et al in 1982 by the name of triacontanol from *B.diffusa* [20]. During 1975-75, Proteins, Carbohydrates and Lipids Glycoprotein (an Antiviral agent) Protein and free Amino acids were reported by different authors [21-24]. Then a lot of compounds have been isolated and reported [4] and the story is still at run.








 Table 1: List of different compounds isolated from different species of *Boerhavia*

S.No.	Name of compound	Mol. Formula	Mol. Wt.	Physical and spectral data [mp °C/[α]D (°)/IR cm ⁻¹ /UV nm]	Source	Ref.
1	Isorhamnetin glycosides			-/-/-	<i>B. erecta</i>	4
	Kaempferol (78)	C ₁₅ H ₁₀ O ₆	286.24	276-278/-/204, 265, 365	<i>B. erecta</i>	4
	Quercetin (77)	C ₁₅ H ₁₀ O ₇	302.24	313-314 dec/-/258, 375	<i>B. erecta</i>	4
2	Catechin (71)	C ₁₅ H ₁₄ O ₆	290.272	93-96/+17/-/-	<i>B. erecta</i>	4
3	Betanin (72)	C ₂₄ H ₂₆ N ₂ O ₁₃	550.475	-/-/-	<i>B. erecta</i>	4
4	Isobetanin (73)	C ₂₄ H ₂₆ N ₂ O ₁₃	550.475		<i>B. erecta</i>	4
5	Neobetanin (74)	C ₂₄ H ₂₄ N ₂ O ₁₃	548.459	-/-/-/267, 306, 470	<i>B. erecta</i>	4
6	Procynidins (75)	[C ₁₇ H ₁₈ O ₆] ₄	1273.28	-/-/-	<i>B. erecta</i>	4
7	5-Methyleicos-4-ene, Eicos-4-ene, 4-Methyloctadec-3-ene and 4-Methylnonadecylbenzene			-/-/-	<i>B. diffusa</i>	5
8	Tannins, Saponins and Calcium oxalate			-/-/-	<i>B. coccinea</i> <i>B. erecta</i> <i>B. diffusa</i>	6
9	Protein, β -Carotene, Ascorbic acid			-/-/-	<i>B. diffusa</i>	7
10	Boerhavisterol (55)	C ₂₉ H ₅₀ O	414.713	128-139/-/3480, 2955, 2880, 1600, 1455, 1340/265 (MeOH)	<i>B. diffusa</i>	8
11	Boerhavisterol acetate (56)	C ₃₁ H ₅₂ O ₂	456.74	81-82/-/1725, 1600/-	<i>B. diffusa</i>	8
12	Boerhadiffusene (57)	C ₃₀ H ₅₄	414.757	103-104/-/2940, 2855, 1625, 1455, 1445/272 (MeOH)	<i>B. diffusa</i>	8
13	Boerhavilanosteryl benzoate (58)	C ₄₃ H ₆₄ O ₈	708.974	250-251dec/-/3400, 2935, 2870, 1725, 1710, 1640, 1585, 1440/217, 280	<i>B. diffusa</i>	8

14	Aglycone (59)	C ₃₀ H ₅₀ O ₃	458.72	202-203/-/-	<i>B.diffusaa</i>	8
15	Lanost-ene-3,27-diol (60)	C ₃₀ H ₅₂ O ₃	460.73	161-162/-/-	<i>B. diffusa</i>	8
16	Diffusarotenoid (61)	C ₂₂ H ₂₀ O ₇	396.396	261-263 (dec.)/-3300, 2965, 1730, 1645, 1615, 1585, 1440 / 218, 264, 291, 355	<i>B. diffusa</i>	8
17	Methyl diffusarotenoid (62)	C ₂₄ H ₂₄ O ₇	424.44	241-242/-/1730, 1710, 1640, 1610/-	<i>B. diffusa</i>	8
18	Coccineone E (51)	C ₁₈ H ₁₆ O ₇	344.32	206-209/-25.5 (CHCl ₃)/ 3366, 2971, 1638, 1580, 1491, 1451/203, 282,	<i>B.coccinea</i>	9
19	Choline (70)	C ₅ H ₁₄ NO ⁺	104.172	-/-/-	<i>B.diffusa</i>	10
20	Eupalitin 3-O-β-D-galactopyranosyl-(1→2)-β-D-glucopyranoside (25)	C ₂₉ H ₃₅ O ₁₇	655.187	-/-19.2 (MeOH)/ 3350, 2900, 2200, 1650, 1590, 1440/-	<i>B.repens</i>	11
21	Eupalitin 3-O-β-D-galactopyranoside (26)	C ₂₃ H ₂₅ O ₁₂	493.135	-/-24.0(MeOH) /3400, 2900, 2200, 1600, 1507, 1440/-	<i>B.repens</i>	11
22	6-methoxykaempferol-3-O-β-D-(1→6)rhamnoside (27)	C ₂₈ H ₃₃ O ₁₆	625.177	-/-30.6 (MeOH) /3400, 2900, 2200, 1600, 1507, 1440/-	<i>B.repens</i>	11
23	Borhavine (9)	C ₂₀ H ₁₈ O ₇	370.358	230-232/-/ 2940, 1660, 1630, 1610, 1590, 1500, 148 0/225, 275, 314, 355	<i>B. diffusa</i>	12
24	Acetylborhavine (10)	C ₂₂ H ₂₀ O ₈	412.389	210-212/-/ 2940, 2820, 1870, 1655, 1620, 1480, 1400, 1380, 1300, 1215/-	<i>B. diffusa</i>	12
25	11-methoxyborhavine (11)	C ₂₁ H ₂₀ O ₇	384.379	220-222/-/-	<i>B. diffusa</i>	12
26	Demethylatedborhavine (12)	C ₁₉ H ₁₆ O ₇	356.326	215-218/-/-	<i>B. diffusa</i>	12
27	Diacetylborhavine (13)	C ₂₀ H ₁₈ O ₇	370.353	214-215/-/-	<i>B. diffusa</i>	12
28	1-Hdroxy-4,5-dimethylxanthone (14)	C ₁₅ H ₁₂ O ₃	240.254	-/-/-	<i>B. diffusa</i>	12
29	Coccineone C (49)	C ₁₇ H ₁₄ O ₇	330.293	-/-339 (MeOH)/-/-	<i>B.coccinea</i>	13
30	Boeravinone D (3)	C ₁₈ H ₁₄ O ₇	342.304	-/0 (acetone)/3350, 1645, 1619, 1597, 1510, 1465/217, 277, 301, 348	<i>B. diffusa</i>	14
31	Boeravinone E (4)	C ₁₇ H ₁₂ O ₇	328.278	-/0 (acetone)/-/217, 278, 300, 349	<i>B. diffusa</i>	14
32	Boeravinone F (5)	C ₁₇ H ₁₀ O ₇	326.262	-/-1721, 1646/217, 265, 294, 330	<i>B. diffusa</i>	14
33	Liriodendrin (21)	C ₃₄ H ₄₆ O ₁₈	742.27	256-259 /-13.5 (pyridine)/3450, 1597, 1510, 1465/272, 281	<i>B. diffusa</i>	15
34	Syringaresinol-β-D-glusoside (22)	C ₂₈ H ₃₆ O ₁₃	580	-/-10 (MeOH) / 3350, 1590, 1510, 1460/270	<i>B. diffusa</i>	15
35	Ursolic acid (66)	C ₃₀ H ₄₈ O ₃	456.36	291/+66 (EtOH+KOH)/-/-	<i>B.diffusaa</i>	15
36	Repenone (15)	C ₁₈ H ₁₂ O ₈	356.288	-/- 3520, 3600, 1730, 1610/-	<i>B.repens</i>	16
37	Repenol (16)	C ₁₈ H ₁₂ O ₉	372.287	-/- 3420, 3520, 1720, 1640, 1610/-	<i>B.repens</i>	16
38	Punarnavoside (29)	C ₂₈ H ₃₀ O ₁₀	526.539	185-186/-/ 3400-	<i>B. diffusa</i>	17

					3350,2900, 1735,1650,1510,1460,142 0,1330,1270,1230,1180,1 160,1120,1070,920,900,8 30/-		
39	Peracetylpunarnavoside (30)	C ₄₀ H ₄₂ O ₁₆	778.25	118-20/-/ 2850, 1740, 1600, 1500,1450/-	<i>B.diffusaa</i>	17	
40	Methyl punarnavoside (31)	C ₃₀ H ₃₄ O ₁₀	554.215	156-57/-/ 3300, 2900, 2850, 1710-1700, 1620, 1525, 1440/-	<i>B.diffusaa</i>	17	
41	p-Hydroxyphenylpropionic acid (32)	C ₁₀ H ₁₂ O ₃	180.08	132-133/-/-	<i>B.diffusaa</i>	17	
42	p-methoxyphenylpropionic acid (33)	C ₉ H ₁₂ O ₃	168.079	90-95/-/-	<i>B.diffusaa</i>	17	
43	Aglycone A (34)	C ₂₆ H ₂₄ O ₇	448.152	88-90/-/3400, 2900, 1735, 41640,1600,1520,1410/-	<i>B.diffusaa</i>	17	
44	Aglycone A acetate (35)	C ₂₈ H ₂₆ O ₈	490.163	110-12/-/2850, 1740- 1735, 1620,1590,1540,1420/-	<i>B.diffusaa</i>	17	
45	Aglycone B (36)	C ₂₄ H ₂₄ O ₅	392.162	102-5/-/3400, 2850, 1725, 1640, 1590, 1520, 1470, 1430/-	<i>B.diffusaa</i>	17	
46	Aglycone B methyl (37)	C ₂₅ H ₂₆ O ₅	406.178	131-132/-/ 2900,1730, 1635, 1590,1540,1570,1490/-	<i>B.diffusaa</i>	17	
47	1-glucopyrano-2-benzyl-4,5-dihydroxybenzene (38)	C ₁₈ H ₂₂ O ₈	366.13	158-159/-/ 3550, 2900, 1600, 1510, 1460, 1420, 1250,1210,1150,1120,930 ,830,780/-	<i>B.diffusaa</i>	17	
48	Aglycone C (39)	C ₁₅ H ₁₆ O ₃	244.110	185-86/-/3450, 1640, 1610, 1570,1550,1470/-	<i>B.diffusaa</i>	17	
49	Aglycone C methyl (40)	C ₁₆ H ₁₈ O ₃	258.126	195-97/-/1640, 1610, 1590, 1550,1510,1495,1485/-	<i>B.diffusaa</i>	17	
50	2,4,5-Trihydroxydiphenyl methane (41)	C ₁₃ H ₁₂ O ₃	216.233	-/-/-	<i>B.diffusaa</i>	17	
51	Boeravinone A (1)	C ₁₈ H ₁₄ O ₆	326.305	215-217/0 (acetone)/3600, 3250, 1650, 1620, 1595, 1495/217, 276, 300, 340	<i>B. diffusa</i>	18	
52	Boeravinone B (2)	C ₁₇ H ₁₂ O ₆	312.278	-/0 (acetone)/3250, 1650, 1620, 1595, 1495/217, 273, 300, 345	<i>B. diffusa</i>	18	
53	Methylarachidate, Methylbehenic acid, Methylbehenate, Methylheptadecyclate, Methyloleate, Methylpalmitate, and Methylstearate			-/-3000, 1710/-	<i>B.diffusaa</i>	18	
54	β-Sitosterol (42)	C ₂₉ H ₅₀ O	414.71	140/-37 (CHCl ₃)/ 3450- 2450, 1650, 815/-	<i>B.diffusaa</i>	18	
55	Stigmasterol (43)	C ₂₉ H ₄₈ O	412.70	170/-51 (CHCl ₃)/ 3400,3015,1615,810/-	<i>B.diffusaa</i>	18	
56	Campesterol (44)	C ₂₈ H ₄₈ O	400.69	157-158/-33 (CHCl ₃)/ 3440,3045,1620, 815/-	<i>B.diffusaa</i>	18	

57	Sitosteryl palmatite (45)	C ₄₅ H ₈₀ O ₂	653.116	-/-/-	<i>B.diffusaa</i>	18
58	Sitosteryl oleate (46)	C ₄₇ H ₈₂ O ₂	679.153	-/-1720, 1669/-	<i>B.diffusaa</i>	18
59	Palmitoyl glucosyl sitosterol (52)	C ₅₁ H ₉₀ O ₇	815.256	-/-47.7(CHCl ₃) /3595-3399, 1733/-	<i>B.diffusaa</i>	18
60	Palmitoylglucosyltriacetate sitosterol (53)	C ₅₇ H ₉₆ O ₇	893.368	-/-/1755/-	<i>B.diffusa</i>	18
61	Palmitoylglucosyltetraacetate sitosterol (54)	C ₄₃ H ₆₈ O ₆	680.996	-/-/-	<i>B.diffusa</i>	18
62	Boeravinone C (6)	C ₁₈ H ₁₆ O ₇	344.320	248-249/- 47(acetone)/3550, 3440, 1630, 1580, 1510, 1480 /331,293,212,207	<i>B. diffusa</i>	19
63	Boerhavinone C diacetate (7)	C ₂₂ H ₂₀ O ₉	428.389	-/-1760/-	<i>B. diffusa</i>	19
64	Boerhavinone C triacetate (8)	C ₂₄ H ₂₂ O ₁₀	470.425	-/-1760,1745/-	<i>B. diffusa</i>	19
65	Alcohol (23)	C ₁₈ H ₁₈ O ₇	346.11	208-209/-/-	<i>B.diffusaa</i>	19
66	Epimeric Alcohol (24)	C ₁₈ H ₁₈ O ₇	346.11	203-205/-/-	<i>B.diffusaa</i>	19
67	Linolenate, Palmitate, and Linoleate			-/-/-	<i>B.diffusa</i> <i>B.erecta</i> <i>B.repens</i>	20
68	5,23-Stigmastadienol and 5,24(25)Stigmastadienol,			-/-/-	<i>B.diffusa</i> <i>B.erecta</i> <i>B.repens</i>	20
69	Stigmastanol (64)	C ₂₉ H ₅₂ O	416.72	136-137/+24.8/-/-	<i>B.diffusa</i> <i>B.erecta</i> <i>B.repens</i>	20
70	Coccineone A (47)	C ₁₅ H ₈ O ₆	284.225	310/-/222, 254, 320	<i>B.coccinea</i>	21
71	Coccineone B (48)	C ₁₆ H ₁₀ O ₆	298.251	>310/recemic/-/214, 268, 300, 330/	<i>B.coccinea</i>	21
72	Coccineone D (50)	C ₁₆ H ₁₂ O ₇	316.267	237-240/-440 (MeOH)/-/-	<i>B.coccinea</i>	21
73	Borhavone / C-Methylflavone (17)	C ₁₉ H ₁₈ O ₆	342.348	222-223/(MeOH) /-/3460, 2900, 1650, 1600, 1200/ 275,314,350	<i>B. diffusa</i>	22
74	Borhavone diacetae (18)	C ₂₃ H ₂₂ O ₈	426.416	-/-1780, 1200/-	<i>B. diffusa</i>	22
75	Demethylated borhavone (19)	C ₁₇ H ₁₄ O ₆	314.289	283-85/-/275, 300, 350 (MeOH)	<i>B. diffusa</i>	22
76	Borhavone tetraacetate (20)	C ₂₅ H ₂₂ O ₉	466.337	166-67/-/-	<i>B. diffusa</i>	22
77	β-Ecdysone (67)	C ₂₇ H ₄₄ O ₇	480.31	240-241/-3500,1650/240	<i>B.diffusaa</i>	23
78	Triacontanol / Myricyl alcohol (69)	C ₃₀ H ₆₂ O	438.819	87/-/3500,1650/240 (MeOH)	<i>B.diffusa</i> <i>B.repens</i>	23 28
79	Proteins, Carbohydrates and Lipids			-/-/-	<i>B.diffusa</i>	24
80	Glycoprotein (an Antiviral agent)		16000- 20000 daltons	-/-/-	<i>B.diffusa</i>	25
81	Protein and free Amino acids			-/-/-	<i>B.diffusa</i>	26

82	D-glucose, D-xylose, D-glucoronic acid, D-galactose, L-arabinose, and L-Rhamnose,		-/-/-	<i>B.repens</i>	27
83	Myristic acid (68)	C ₉ H ₈ O ₅	196.037	212/-/-	<i>B.repens</i> 28
84	Hentriacontane (65)	C ₃₁ H ₆₄	436.847	65.9/-/-	<i>B.diffusaa</i> 29
85	Hypoxanthine-9-L-arabinofuranoside (28)	C ₁₁ H ₁₃ N ₃ O ₅	267.24	-/-/-	<i>B.diffusa</i> 30
86	Punarnavine	C ₁₇ H ₂₂ N ₂ O	270.374	236-237dec./-/-	<i>B. diffusa</i> 31, 34
87	(Z)-24-Triacontenoic acid (63)	C ₃₀ H ₅₈ O ₂	450.787	60.8-61.2/-/-	<i>B.repens</i> 32
88	Alanine Arachidic Acid Aspartic Acid Behenic Acid Virus Inhibitor Boerhaavic Acid Dauosterol 5-7-dihydroxy-3'-4'-dimethoxy-6-8-dimethyl Fructose Glutamic Acid Glutamine Glycerol Glycine Heptadecyclic Acid Histidine Leucine Methionine Oleaic Acid Oxalic Acid Palmitic Acid Proline Proline, hydroxy Serine Stearic Acid Sucrose Threonine Tyrosine Valine		-/-/-	<i>B. diffusa</i> 33	

References:

1. Watt, G., *A Dictionary of Economic Products of India*, **1972**, vol. 1, pp485-486.
2. *Indian Medicinal Plants*, third enlarged edition, vol. 9, pp2818-19.
3. Nasir, E.; Ali, E.I.; *Flora of West Pakistan*, Fakhri Printing Press Karachi **1977**, vol. 115: pp1-16.
4. Stintzing, F.C.; Kammerer, D.; Schieber, A.; Adama, H.; Nacoulma, O.G.; Carle, R., *J. Biosciences*, **2004**, 59(1/2), 1-8.
5. Singh, B.L.; Verma, R.K.; Singh, S.C.; Gupta, M.M., *J. Indian chemical society*, **2002**, 79(11), 911-912.
6. Edeoga, H.O.; Ikem, C.I.; *South African Journal of Botany*, **2002**, 68(3), 382-385.
7. Bharati, R.; Umameheshwari, K.; *Journal of Food Science and Technology*, **2001**, 38(5), 534-536.
8. Gupta J.; Ali, M., *Indian J. Chem., Sect. B*, **1998**, 37, 912-917.
9. Santos, A.S.; Caetano, L.C.; Sant'Ana, A.E.G., *Phytochemistry*, **1998**, 49(1), 255-258.
10. Wahi, A.K.; Agarwal, U.K.; Gupta, R.C., *Natl. Acad. Sci. Lett. (India)*, **1997**, 20(9&10), 119-123.
11. Li, J.; Li, H.; Kadota, S.; Numba, T., *J. Nat. Prod.*, **1996**, 59, 1015-1018.
12. Ahmed, B. Yu, C. P., *Phytochemistry*, **1992**, 31(12), 4382-4.
13. Ferrari, F.; Messana, I., *J. Nat. Prod.*, **1991**, 54(2), 597-598.
14. Lami, N.; Kadota, S.; Kikuchi, T., *Chem. Pharm. Bull.*, **1991**, 39(7) 1863-1865.
15. Lami, N.; Kadota, S.; Kikuchi, T.; Momose, Y., *Chem. Pharm. Bull.*, **1991**, 39(6), 1551-1555.
16. Ahmed, N.; Dhatta, B.K.; Rauf, S.S., *Phytochemistry*, **1990**, 29(5), 1709-10.
17. Jain, G.K.; Khanna, N.M., *Indian J. Chem., Sect. B*, **1989**, 28, 163-166.
18. Shigetoshi, K.; Nzunzu, L.N.; Yasuhiro, T.; Tohru, K., *Chem. Pharm. Bull.*, **1989**, 37(12), 3214-3220.
19. Katoda, S.K.; Nzunzu, L.; Tezuka, Y.; Kikuchi, T., *Chem. Pharm. Bull.*, **1988**, 36(6), 2289-2292.
20. Miralles, J.; Noba, K.; Ba, T. A.; Gaydou, M. E.; Kornprobst, J.M., *Biochem. Syst. Ecol.*, **1988**, 16(5), 475-478.
21. Messana, I.; Ferrari, F.; Goulart, A.E.; Sant'Ana, *Phytochemistry*, **1986**, 25(11), 2688-89.
22. Gupta, D.R.; Ahmad, B., *Indian J. Chem., Sect. B*, **1984**, 23, 682-684.
23. Suri, O.P.; Kant, R.; Jamwal, R.S.; Suri, K.A.; Atal, C.K., *Planta Medica*, **1982**, 44(3), 180-181.
24. Vasi, I.G.; Kalintha, V.P.; *J. Indian Chem.* **1979**, 51(5), 214-216.
25. Verma, H.N.; Awasthi, L.P.; Saxena, V., *Can. J. Bot.*, **1979**, 57(11), 1214-1217.
26. Goerge, S.P.; Mehta, R.; Jain, V.M.; Chaudhry, M.R.; Bhatnagar, R., *Trans. Indian Soc. Desert Technol. Univ. Cent. Desert Study*, **1978**, 3(1), 31-34.
27. Haq, Q.; Haq, N., *Bangladesh J. Sci. Ind. Res.* **1975**, 10(1-2), 84-7.

28. Khalique, A.; Hannun, A.; Rehman, J.; Kiamuddin, M., *Bangladesh J. Sci. Ind. Res.*, **1974**, 9(1-2), 47-50.
29. Misra, A.N.; Tiwari, H.P., *Photochemistry*, **1971**, 10(12), 3318-19.
30. Kamaluddin, A.; Afzal, H., *Pak. J. Biol. Agr. Sci.*, **1968**, 11(2), 41-44.
31. Bhalla T.N.; Gupta, M.L., *J. Res. Indian*, **1968**, 12, 37.
32. Lightelm. S. P., *J. Sci. Food Agric.*, **1954**, 5, 281.
33. © Copyrighted 2004. Raintree Nutrition, Inc. Carson City, NV 89701. All Rights Reserved. www.rain-tree.com
34. Basu. N.K., *Q.J. Pharm. Pharmacol.*, **1947**, 20, 41.
35. Ujowundu1, C.O.; Igwe1, C.U.; Enemor, V.H.A.; Nwaogu, L.A.; Okafor, O.E., *Pakistan Journal of Nutrition*, **2008**, 7 (1): 90-92, ISSN 1680-5194.
36. Chude, M.A.; Orisakwe, O.E.; Afonne, O.J; Gamaniel, K.S.; Vongtau, O.H.; Obi, E., *Indian Journal of Pharmacology*, **2001**, 33, 215-216.